RESEARCH

Open Access

Anticancer and cancer preventive activities of shogaol and curcumin from Zingiberaceae family plants in KG-1a leukemic stem cells



Pawaret Panyajai¹, Natsima Viriyaadhammaa¹, Sawitree Chiampanichayakul^{1,2,3}, Yasuhisa Sakamoto⁴, Siriporn Okonogi^{3,5}, Toshiro Moroishi^{4,6*} and Songyot Anuchapreeda^{1,2,3*}

Abstract

Background Leukemic stem cells (LSCs) present a significant challenge in the treatment of leukemia in patients because they exhibit a drug-resistant phenotype, making them difficult to eliminate. Searching for a new anticancer drug is crucial for improving leukemia treatment. Plants from the Zingiberaceae family are frequently used in traditional medicines due to their safety and accessibility. This study explores the anticancer activity, cancer preventive properties, and apoptosis inducing mechanisms of active compounds derived from these plants.

Methods Ten crude ethanolic extracts from each plant of the Zingiberaceae family were obtained using maceration techniques. The cytotoxicity of all extracts anticancer was assessed in comparison to anticancer drugs (cyclophosphamide, cytarabine, doxorubicin, and idarubicin) using MTT assay on cancer cell lines (KG-1a, K562, A549, MCF-7, and HeLa) and peripheral blood mononuclear cells (PBMCs). Cancer prevention properties of the effective extracts and their active compounds were evaluated by measuring the levels of tumor necrosis factor-alpha (TNF-a), interleukin-2 (IL-2), and nitric oxide (NO) using commercial kits. Cell cycle and cell death analyses were conducted using flow cytometry. Moreover, the effects of effective extracts and their active compounds on WT1 and CD34 expressions, as well as the apoptosis mechanism induced by the active compounds in KG-1a cells, were determined by Western blotting.

Results The cytotoxicity tests revealed that crude ethanolic extracts from *Curcuma longa, C. zedoaria*, and *Zingiber officinale* exhibited effective cytotoxicity against cancer cell lines while demonstrating lower impact on PBMCs. The active compounds of *C. longa* and *C. zedoaria* are curcuminoids, while those in *Z. officinale* are shogaol and gingerol. Notably, the IC_{20} values of curcuminoids and shogaol exhibited cancer prevention properties and reduced WT1 protein expression, thereby inhibiting cell proliferation. Furthermore, shogaol and curcumin demonstrated the ability to arrest the cell cycle at the G_2/M phase and induce apoptosis through the Akt pathway.

Conclusion These findings highlight shogaol and curcumin as promising compounds for leukemia treatment.

*Correspondence: Toshiro Moroishi moroishi@kumamoto-u.ac.jp Songyot Anuchapreeda songyot.anuch@cmu.ac.th

Full list of author information is available at the end of the article



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by-nc-nd/4.0/.

Keywords Zingiberaceae family, Cancer, Leukemia, Leukemia stem cell, Curcumin, Shogaol, WT1, Cancer prevention property, Apoptosis

Introduction

Chemotherapy is one of the effective methods for leukemia treatments. However, most of the chemotherapies function by obstructing DNA replication, a mechanism that affects both cancer and normal cells alike, leading to unpleasant side effects for patients. Plants from the Zingiberaceae family are commonly used in traditional medicines and as cooking ingredients in Southeast Asia. Most plants in this family are extensively employed in traditional remedies to address issues such as stomachaches, hemorrhoids, blood circulation, and muscular pain relief. They are also used in herbal compress balls for massages. Moreover, rhizomes and leaves of these plants are used as spices and cooking ingredients. Previous reports have highlighted the diverse biological activities from this family [1–4], presenting an intriguing avenue for developing anticancer drugs with reduced side effects, improved accessibility, and enhanced safety.

Unregulated inflammation is often associated with the development and progression of cancer since inflammatory response can damage the DNA of normal cells. This damage increases the risk of mutations that eventually contribute to cancer [5-7]. Several active compounds from plants, such as sulforaphane, lycopene, curcumin, shogaol, and resveratrol, exhibited cancer prevention properties by suppressing the production of inflammatory cytokines and nitric oxide (NO), while enhancing the phagocytic activity of macrophages [8-12]. Thus, suppression of unregulated inflammation represents a potential strategy for preventing cancer initiation [13].

Currently, numerous studies have highlighted leukemic stem cells (LSCs) as a significant challenge in patients with leukemia. These cells exhibit surface marker phenotypes CD34⁺/CD38⁻ and share characteristics with hematopoietic stem cells (HSCs) [14, 15]. Additionally, they demonstrated greater resistance against chemotherapeutic drugs compared to normal leukemic cells, making their elimination crucial for improving treatment outcomes. Wilms' tumor 1 (WT1) plays an active role in cell growth and development during normal hematopoiesis [16–20]. However, the overexpression of WT1 protein can induce leukemic cell proliferation. Consequently, the overexpression of the WT1 gene has been used as a biological marker for diagnosing and evaluating minimal residual disease (MRD) in leukemia [21]. Furthermore, the elevated expression of WT1 protein is associated with poor prognosis and worse long-term outcomes in acute myeloid leukemia (AML) [21-23], often leading to relapse [24]. These findings suggested that WT1 plays a crucial role in leukemogenesis, as it is present in both leukemia cells and leukemic stem cells. Simultaneously, many studies have revealed the crucial role of the Hippo signaling pathway in regulating organ size, tissue homeostasis, and cell proliferation. Dysregulation of this pathway also contributes to various diseases, including cancer [25, 26]. In leukemia, Yes-associated protein (YAP) plays important roles in proliferation and leukemogenesis, particularly in chronic myeloid leukemia (CML) [27]. YAP is a key molecule in Hippo signaling pathway. Furthermore, inhibiting YAP function has been shown to limit cell proliferation and induce apoptosis in HL-60 cells [28]. Thus, targeting the Hippo signaling pathway may offer novel opportunities for overcoming resistance in leukemic stem cells and advancing leukemia treatment.

While numerous reports have explored the biological activities of compounds of plants from the Zingiberaceae family, there is less information regarding the effects of extracts or active compounds from this family on WT1 protein expression and the Hippo signaling pathway in LSCs. To investigate this unexplored aspect, this study aims to evaluate the cytotoxicity of ten plant extracts from the Zingiberaceae family in a leukemic stem cell line, comparing their effects to those on other cancer cell lines and peripheral blood mononuclear cells (PBMCs). Additionally, we aim to explore the cancer preventive properties of extracts from candidate plants and investigate the apoptotic mechanisms induced by active compounds in KG-1a leukemic stem cell line. By assessing their impact on pathways associated with leukemogenesis, this study aims to provide new insights into the therapeutic potential of these plants for leukemia treatment.

Materials and methods

Crude ethanolic extraction

In June 2019, fresh rhizomes of Alpinia galanga, Boesenbergia rotunda, Curcuma aeruginosa, C. longa, C. mangga, C. zedoaria, Kaempferia parviflora, Zingiber montanum, Z. officinale, and Z. ottensii were collected from a local garden in Chiang Mai, Thailand.

The authors confirm that all methods involving the plants, and their materials complied with relevant institutional, national, and international guidelines and legislation. Additionally, these plants are common medicinal plants in Thailand and are widely used in Thai traditional medicine. They are not at risk of extinction. All plant materials were identified and authenticated by Wannaree Charoensup, Chiang Mai University, Chiang Mai, Thailand. Their voucher specimens (deposition no. 009245 for A. galanga, 009724 for B. rotunda, 0023261 for C. aeruginosa, 023356 for C. longa, 0023362 for C.

mangga, 0023369 for *C. zedoaria*, 0023370 for *K. parviflora*, 004581 for *Z. montanum*, 0023361 for *Z. officinale*, 000109 for *Z. ottensii*) were deposited in the Herbarium, the Faculty of Pharmacy, Chiang Mai University, Thailand.

Plant collection was done under the license of the Ghana Forestry Commission, according to the guidelines of the IUCN Policy Statement on Research Involving Species at Risk of Extinction and the Convention on the Trade in Endangered Species of Wild Fauna and Flora.

The ethanolic extracts of the fresh rhizomes were obtained using a modified maceration technique based on a previous study [29]. Briefly, each rhizome was dried and ground into a powder. Then, the powder was macerated with 95% ethanol at a ratio of 1:3 for 48 h. The filtrates were collected by filtration technique using a filter cloth, and the residue was subjected to maceration with 95% ethanol at a ratio of 1:3 for an additional 48 h, repeated three times. The filtrates were evaporated to dryness using a rotary evaporator, and the resulting extracts were stored in light-resistant containers within a -20 °C freezer.

Active compounds

Standard curcuminoids (80% pure curcumin, 17% demethoxycurcumin, and 3% bisdemethoxycurcumin), shogaol, and gingerol were purchased from Sigma-Aldrich (MA, USA). The "in-house" curcuminoids refer to compounds purified in our laboratory. These compounds were extracted from natural sources such as the rhizomes of *C. longa* and *C. zedoaria*, subsequently purified in our laboratory using column chromatography. The purified compounds were then analyzed using HPLC to confirm their identity and purity, as described previously [30, 31]. The in-house curcumin, demethoxycurcumin, and bisdemethoxycurcumin, after column chromatography purification, exhibited 100% purity in each compound, as demonstrated in a previous study [32].

Cell culture

Five cancer cell lines (KG-1a, K562, A549, MCF-7, and HeLa) were used as human cancer cell line models in this study. KG-1a cells were cultured in IMDM (Iscove's Modified Dulbecco's Medium) medium (Invitrogen[™], MA, USA) supplemented with 20% fetal bovine serum (Capricorn Scientific, Ebsdorfergrund, Germany), 100 units/mL penicillin and 100 µg/mL streptomycin (Invitrogen[™], MA, USA). K562 cells were cultured in RPMI (Roswell Park Memorial Institute)-1640 medium (Invitrogen[™], MA, USA) supplemented with 10% fetal bovine serum, 2 mM L-glutamine (Invitrogen[™], MA, USA), 100 units/mL penicillin and 100 µg/mL streptomycin. A549, MCF-7, HeLa, and RAW264.7 cells were cultured in DMEM (Dulbecco's Modified Eagle Medium) medium

(Invitrogen[™], MA, USA) supplemented with 10% fetal bovine serum 100 units/mL penicillin and 100 μ g/mL streptomycin (Invitrogen[™], MA, USA). All cancer cell lines were cultured at 37 °C in a humidified incubator with 5% CO₂.

Isolation of PBMCs

PBMCs were isolated from blood samples using the Ficoll-Hypaque technique. Initially, 10–20 mL of blood was collected in a heparin tube at a ratio of 1:500 from a minimum of three healthy volunteers. Subsequently, the blood samples were diluted with Phosphate buffer saline (PBS), pH 7.4, at a ratio of 1:1 to reduce viscosity. Histopaque-1077 (Sigma-Aldrich, MA, USA) was then underlaid the diluted sample and centrifuged at 400×g for 30 min. The resulting PBMCs were collected, washed with PBS, pH 7.4, and prepared for further experiments.

MTT test

MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay was used to investigate the cytotoxicity of the test samples. KG-1a $(1.5 \times 10^4 \text{ cells/well})$, K562 (1.0×10^4 cells/well), PBMCs (1.0×10^5 cells/well), A549 (5.0×10^3 cells/well), MCF-7 (5.0×10^3 cells/well), and HeLa $(5.0 \times 10^3 \text{ cells/well})$ were seeded into a 96-well plate and incubated overnight at 37 °C with 5% CO₂. Subsequently, the cells were treated with various concentrations of crude ethanolic extract ($0-100 \mu g/mL$), active compounds (0–100 μ g/mL), or chemotherapeutic drugs as positive controls: cyclophosphamide ($0-400 \mu g/mL$), cytarabine (0-100 µg/mL), doxorubicin (0-1,000 ng/ mL), and idarubicin (0-1,000 ng/mL), with 0.4% DMSO as the vehicle control (VC), for 48 h. After that, 5 mg/ mL of MTT dye solution (Sigma-Aldrich, MA, USA) was added and incubated for 4 h. The produced formazan crystals were dissolved in 200 µL of DMSO (Sigma-Aldrich, MA, USA); then, the optical density (OD) was measured using an ELISA plate reader (Metertech, Taipei, Taiwan) at 578 nm with a reference wavelength of 630 nm. The percentage of surviving cells was calculated from the absorbance values of the test and control wells using the following Equation.

% Cell viability =
$$\frac{\text{OD}_{\text{sample}}}{\text{OD}_{\text{vehicle control}}} \times 100$$
 (1)

where OD_{sample} is a mean absorbance in test well and $OD_{vehicle\ control}$ is a mean absorbance in VC well. The average percentage of surviving cells at each concentration obtained from triplicate experiments was plotted as a dose-response curve. Inhibitory concentration values at 50% growth (IC₅₀ value) were defined as the lowest concentration of the test sample that inhibited cell growth by 50% compared to the untreated control.

To calculate IC_{50} values, a dose-response curve was fitted using non-linear regression analysis, as described in the GraphPad Prism guidelines (https://www.graphpad.com/support/faq/how-to-determine-an-icsub50sub/). The IC_{50} value was obtained by determining the point at which the curve intersected 50% growth inhibition. Non-linear regression graphs of the dose-response curve for each compound are provided in the supplementary data file (Figs. S1–S8).

The selectivity index (SI) of active compounds and chemotherapeutic drugs was calculated for using the following Eq.

Selectivity index (SI) =
$$\frac{\text{IC}_{50 \text{ N}}}{\text{IC}_{50 \text{ C}}}$$
 (2)

where $IC_{50 N}$ is IC_{50} for normal cells and $IC_{50 C}$ is IC_{50} for cancer cells which were treated with the same compounds in both cells.

Trypan blue exclusion test

This test was used to confirm the cytotoxicity of the test samples. The cancer cells after exposure to crude ethanolic extracts and active compounds for 48 h were harvested and washed with ice-cold PBS, pH 7.4 for 3 times. Then, the cells were resuspended with PBS, pH 7.4. The cell suspensions were diluted with PBS, pH 7.4 at the appropriate dilution before being mixed with 0.2% trypan blue solution at 1:2 dilution for cell count on a hemocytometer. From this test, the viable cells and the dead cells could also be obviously detected.

Cancer prevention properties

For the measurement of tumor necrosis factor-alpha (TNF- α) and NO, RAW264.7 cells were pre-treated with plant extracts and active compounds at a non-cytotoxic dose (IC₂₀) (Table S9) or 1 μ g/mL of dexamethasone (positive control) in a 24-well plate for 2 h. Subsequently, 1 μg/mL of lipopolysaccharide (LPS) (Cat No. 00497693, Invitrogen[™], MA, USA) was added to each well, and the cells were incubated for an additional 24 h to allow sufficient time for the compounds to exert measurable anti-inflammatory effects while maintaining cell viability. Afterward, supernatants were collected, and the levels of TNF- α and NO were measured using the Mouse TNF alpha Uncoated ELISA Kit (Invitrogen[™], Thermo-Fisher Scientific, MA, USA) and the Total Nitric Oxide Assay Kit (Invitrogen[™], ThermoFisher Scientific, MA, USA), respectively, according to the manufacturer's instructions.

For the measurement of interleukin-2 (IL-2), human PBMCs were pre-treated with plant extracts and active compounds at the IC_{20} concentration (Table S9) or 1 µg/mL of dexamethasone (positive control) in a 24-well plate

for 2 h. Subsequently, 20 μL/mL of phytohemagglutinin (PHA) (Cat. No. 10576015, Thermo Fisher Scientific, MA, USA) was added to each well, and the cells were incubated for an additional 24 h. After that, supernatants were collected, and the levels of IL-2 were measured by the Human IL-2 Uncoated ELISA Kit (Invitrogen[™], ThermoFisher Scientific, MA, USA), according to the manufacturer's instructions.

The percentage of NO or cytokine inhibitions were calculated using the following equation:

% NO or cytokine inhibition =
$$\frac{A - B}{A} \times 100$$
 (3)

where A is the level of cytokine or NO without treatment, and B is the level of cytokine or NO with treatment.

Cell cycle and apoptosis assay

After treating KG-1a cells with active compounds at various concentrations for 48 h, the cells were pulsed with 10 µM BrdU (Sigma-Aldrich, MA, USA) for 20 min at 37 °C in an incubator with 5% CO₂. Subsequently, the cells were harvested, washed with ice-cold PBS, pH 7.4 three times, and then fixed with 70% ethanol in PBS, pH 7.4 at -30 °C. Following fixation, the cells were washed and resuspended in 2 N HCl/0.5% Triton X-100 for 30 min at room temperature (RT). Then, 0.1 M sodium tetraborate decahydrate was added for 2 min at RT to neutralize the acid. Subsequently, the cells were stained with FITC-anti-BrdU antibody for 30 min at RT. After washing, the cells were resuspended in propidium iodide (PI) (Dojindo, Kumamoto, Japan) in PBS, pH 7.4 at a ratio of 1:1,000. The cell cycle phases were detected using a flow cytometer (CytoFLEX S, Beckman Coulter, CA, USA) and analyzed with the CytExpert program (Beckman Coulter, CA, USA).

The flow cytometry data were analyzed to assess both apoptosis and cell cycle distribution. Apoptosis was evaluated by determining the percentage of cells in the sub- G_1 phase using the Nicoletti assay. For cell cycle analysis, populations in the G_1 , S, and G_2/M phases were calculated after excluding the sub- G_1 population to avoid interference from apoptotic cells, allowing for a clearer assessment of cell cycle arrest in different phases. Data were presented in separate graphs for the sub- G_1 population and cell cycle phase distributions.

Cell death analysis

This test was used to confirm the cytotoxicity of crude ethanolic extracts and active compounds. KG-1a cells after exposure to crude ethanolic extracts and active compounds for 48 h were harvested and washed with ice-cold PBS, pH 7.4 for 3 times. Then, the cells were resuspended with PBS, pH 7.4. The cell suspensions were diluted with PBS, pH 7.4 at the appropriate dilution before being mixed with 0.2% trypan blue solution at 1:2 dilution for cell count on a hemocytometer. From this test, the viable cells and the dead cells could also be obviously detected.

Additionally, PI staining was used to investigate the cytotoxicity of the active compounds. KG-1a cells after exposure to the active compounds with various concentration were harvested at different incubation times and washed with ice-cold PBS, pH 7.4 for 2 times. Then, the cells were resuspended with PI in PBS, pH 7.4 at a ratio of 1:10,000. The percentage of dead cells was detected by flow cytometry (CytoFLEX S, Beckman Coulter, CA, USA) and analyzed by CytExpert program (Beckman Coulter, CA, USA).

Both tests provided similar results regarding cell viability and cell death analysis.

Western blotting

KG-1a cells $(1.5 \times 10^5 \text{ cells/mL})$ were treated with compounds or extracts at different incubation times. Then, the cells were harvested and the whole proteins were extracted using a RIPA buffer (25 mM Tris-base, pH 7.6, 0.1% SDS, 1% Triton X-100, 150 mM NaCl, and 1 mM EDTA). The protein concentration was measured using the Folin-Lowry or Bradford assay. The whole protein lysates (30 µg/lane) were then separated through 7.5% SDS-PAGE under reducing conditions and then transferred to PVDF membranes. The membranes were blocked in 5% skim milk in PBS, pH 7.4 at RT for 2 h, then, target proteins were probed with specific primary antibodies (anti-WT1, anti-CD34, anti-GAPDH, anticleaved caspase-3, anti-cleaved PARP, anti-phospho-Akt, anti-phospho-c-Jun, anti-phospho-SAPK/JNK, and anti-YAP/TAZ) indicated in Table S10 which diluted at a ratio of 1:1,000 in PBS, pH 7.4 at RT for 2 h. After incubation, the membranes were rinsed 6 times with 0.1% Tween-20 (Sigma-Aldrich, MA, USA) in PBS, pH 7.4 (PBS-T), each time for 5 min. The reaction was followed by HRP-conjugated secondary antibody at 1:20,000 dilution in PBS, pH 7.4 at RT for 2 h, then, the membranes were rinsed 6 times for 5 min each with PBS-T. The proteins were shown using Immobilon Forte Western HRP substrate (Millipore, MA, USA). Finally, the protein band signal was quantified by using a scan densitometer (Bio-Rad Laboratories, CA, USA).

Statistical analysis

All experiments were performed in triplicate except the apoptosis mechanism in KG-1a cells. The average of triplicate experiments and standard derivation (SD) were used for quantification. The levels of cell populations were compared to VC in each experiment. The results are shown as mean \pm S.D. The SPSS statistics software ver. 22 (SPSS Inc., IL, USA) was used for statistical analysis. Differences between the means of each sample were analyzed by one-way analysis of variance (one-way ANOVA), followed by LSD post-hoc analysis. Statistical significance was considered at p < 0.05.

Results

Cytotoxicity of crude ethanolic extracts on cancer cell lines and PBMCs

The results indicated that each crude ethanolic extract exhibited various levels of cytotoxicity across different cancer cell lines and PBMCs. Notably, the crude ethanolic extracts of C. longa, C. zedoaria, and Z. officinale demonstrated good cytotoxicity in KG-1a cells, with the IC_{50} values of 24.59 ± 1.40 , 24.72 ± 3.39 , and $21.70 \pm 1.83 \ \mu g/$ mL, respectively. According to National Cancer Institute (NCI) criteria, plants with an IC₅₀ of \leq 30 µg/mL are considered promising candidates for further investigation in cancer treatment [33, 34]. Moreover, these three extracts also exhibited substantial cytotoxicity in other cancer cell lines, except A549, while demonstrating low cytotoxicity in PBMCs (Table 1). Raw data for the IC_{50} values, obtained from non-linear regression graphs performed in triplicate, are provided in the supplementary file (Figs S1–S6 and Tables S1–S6).

Table 1 IC₅₀ values of crude ethanolic extracts from Zingiberaceae plants on cancer cell lines and PBMCs after incubation for 48 h

Crude ethanolic extracts	IC ₅₀ (μg/mL)						
	KG-1a	PBMCs	K562	A549	HeLa	MCF-7	
A. galanga	>100	42.11±3.84	49.10±6.13	90.24 ± 3.75	52.51 ± 6.66	46.34 ± 3.02	
B. rotunda	38.46 ± 0.20	39.83 ± 2.27	27.83 ± 4.49	51.60 ± 5.54	30.08 ± 5.52	38.68 ± 5.79	
C. aeruginosa	74.80 ± 2.62	79.11 ± 3.92	64.70 ± 3.63	>100	57.57 ± 8.16	66.05 ± 2.95	
C. longa	24.59 ± 1.40	76.30 ± 9.13	26.32 ± 2.76	80.10 ± 9.23	20.34 ± 0.39	19.56 ± 3.08	
C. mangga	83.97 ± 2.04	>100	48.76 ± 1.58	>100	66.55 ± 1.45	64.32 ± 5.76	
C. zedoaria	24.72 ± 3.39	42.26 ± 3.15	41.17 ± 3.33	>100	24.68 ± 1.91	18.83 ± 2.41	
K. parviflora	29.34 ± 5.09	46.77 ± 1.48	52.38 ± 7.18	71.33 ± 2.57	15.88 ± 0.97	24.48 ± 1.17	
Z. montanum	46.65 ± 1.82	56.25 ± 3.41	66.14 ± 3.79	>100	58.92 ± 5.15	36.05 ± 6.36	
Z. officinale	21.70±1.83	69.38 ± 10.50	37.76 ± 5.40	86.37 ± 6.81	12.38 ± 1.49	11.78 ± 1.73	
Z. ottensii	50.66 ± 8.14	> 100	>100	>100	> 100	26.69 ± 8.35	

Table 1 IC_{50} values of crude ethanolic extracts from Zingiberaceae plants on cancer cell lines and PBMCs after incubation for 48 h.

Cytotoxicity of active compounds in KG-1a cells and PBMCs

The active compounds of C. longa and C. zedoaria are curcuminoids, including curcumin, demethoxycurcumin, and bisdemethoxycurcumin, while Z. officinale contains shogaol and gingerol. The purity of all curcuminoids was verified using HPLC, as reported in previous studies [30, 31]. HPLC analysis confirmed that in-house curcumin, demethoxycurcumin, and bisdemethoxycurcumin achieved 100% purity, as showed in a previous study [32]. Cytotoxicity analysis using MTT assay indicated that shogaol and three curcuminoids (curcumin, demethoxycurcumin, and bisdemethoxycurcumin) demonstrated significant cytotoxicity against KG-1a cells, with IC_{50} values of 2.99 ± 0.01 , 10.52 ± 0.12 , 11.28 ± 0.85 , and $17.91 \pm 0.99 \ \mu g/mL$, respectively. Furthermore, all active compounds exhibited higher efficacy in cytotoxicity compared to cytarabine and cyclophosphamide against KG-1a cells. When evaluating cytotoxicity in PBMCs, all active compounds and chemotherapeutic drugs (cvclophosphamide, cytarabine, doxorubicin, and idarubicin) demonstrated lower cytotoxicity in PBMCs than in KG-1a cells, except for bisdemethoxycurcumin (Table 2). Raw data for the IC₅₀ values, obtained from non-linear regression graphs performed in triplicate, are provided in the supplementary data (Figs S7–S8 and Tables S7–S8).

SI represents the ratio of the toxic concentration to the effective bioactive concentration of a sample. Ideally, compounds or drugs with a high SI value have a relatively high toxic concentration and a very low active concentration [35, 36]. In this study, the SI calculation revealed that shogaol, gingerol, curcumin, demethoxycurcumin, doxorubicin, and idarubicin had index values greater than 1, indicating that these compounds and drugs exhibited

Table 2 IC₅₀ values and SI of active compounds and chemotherapeutic drugs in KG-1a cells and PBMCs after incubation for 48 h

Compounds and drugs	IC ₅₀	Selec-		
	KG-1a	PBMCs	tivity index (SI)	
Shogaol (µg/mL)	2.99±0.01	9.18±0.82	3.07	
Gingerol (µg/mL)	76.77 ± 1.28	>100	>1.30	
Curcumin (µg/mL)	10.52 ± 0.12	13.78±0.82	1.31	
Demethoxycurcumin (µg/mL)	11.28 ± 0.85	13.87±0.28	1.23	
Bisdemethoxycurcumin (µg/ mL)	17.91±0.99	11.64±1.25	0.65	
Cyclophosphamide (µg/mL)	>400	>400	>1	
Cytarabine (µg/mL)	>100	>100	>1	
Doxorubicin (ng/mL)	652.18 ± 142.06	> 1,000	> 1.53	
Idarubicin (ng/mL)	31.27 ± 5.04	> 1,000	> 31.98	

greater specificity towards KG-1a cells compared to normal PBMCs (Table 2).

Cancer prevention properties and cytotoxic effects of crude ethanolic extracts and their active compounds in cancer cell line and PBMCs

To investigate cancer prevention properties of plant extracts compared to their active compounds, NO and pro-inflammatory cytokines (IL-2 and TNF- α) were investigated in RAW264.7 cells and normal PBMCs after inflammatory induction and treated with IC₂₀ value of each extract and active compound (Table S9). A noncytotoxic dose is used to avoid cell death when determining target proteins in live cells. The result showed that all compounds did not show cytotoxicity on RAW264.7 and PBMCs after incubation for 24 h (Figs. S10 and S11). Each compound and extract had a different inhibitory effect on NO and cytokines.

Compared to dexamethasone (positive control), all compounds and extracts inhibited NO release by more than 20% compared to positive control, except crude ethanolic extract of *Z. officinale* and gingerol. When comparing the activity of the crude ethanolic extract to its active compounds, gingerol showed no significant difference from the crude ethanolic extract of *Z. officinale*. In contrast, shogaol exhibited significantly higher activity, inhibiting NO by $46.46 \pm 3.71\%$, which was notably greater than that of gingerol and its crude ethanolic extract, and even more effective than dexamethasone. Regarding *C. longa*, and *C. zedoaria*, all curcuminoids and their crude ethanolic extract showed no significant difference in NO inhibitory activity, inhibiting approximately 35% (Fig. 1).

While most compounds and extracts inhibited TNF- α release by more than 20%, all showed significantly lower activity than the positive control. However, only gingerol showed less than 10% inhibitory activity. When comparing the activity of the crude ethanolic extract to their active compounds, shogaol exhibited higher inhibitory activity (24.41 ± 0.99%) compared to the crude ethanolic extract of *Z. officinale* and gingerol. Regarding *C. longa* and *C. zedoaria*, all curcuminoids and their crude ethanolic extracts showed no significant difference in TNF- α inhibitory activity with inhibition of approximately 25% or more (Fig. 2).

All compounds and extracts also demonstrated effective inhibitory activity for IL-2 release when compared to positive control, with more than 20% inhibitory activity. However, while most crude ethanolic extracts and active compounds exhibited significantly higher inhibitory activity than positive control, gingerol showed significantly lower inhibitory activity. Comparing the activity between crude ethanolic extract and their active compounds, shogaol exhibited a significantly higher



Fig. 1 Inhibitory effects of crude ethanolic extracts and their active compounds on NO production. LPS-activated RAW264.7 cells were incubated with crude ethanolic extracts and their active compounds for 24 h. The supernatants were collected and measured the levels of NO were measured by NO assay. Shogaol, crude ethanolic extracts of *C. longa* and *C. zedoaria*, and all curcuminoids significantly reduced NO production compared to dexamethasone (positive control), while the crude extract of *Z. officinale* and gingerol had no effect. Data are the mean \pm SD (n=3); *p < 0.05 vs. positive control



Fig. 2 Inhibitory effects of crude ethanolic extracts and their active compound on TNF- α production. LPS-activated RAW264.7 cells were incubated with crude ethanolic extracts and their active compounds for 24 h. The supernatants were collected and measured the levels of TNF- α by ELISA kit. Most compounds and crude ethanolic extracts inhibited TNF- α release by more than 20%, with the crude ethanolic extract of *Z*. officinale and gingerol showing less than 20% inhibition. All treatments showed significantly lower activity than dexamethasone (positive control). Data are the mean ± SD (*n*=3); **p* < 0.05 vs. positive control

inhibitory activity by $80.04 \pm 4.82\%$ when compared to the crude ethanolic extract of *Z. officinale* and gingerol. For *C. longa* and *C. zedoaria*, all curcuminoids and their crude ethanolic extracts showed no significant difference in TNF- α inhibitory activity, which was suppressed by approximately more than 60% (Fig. 3).

Effect of crude ethanolic extracts and active compounds on WT1 and CD34 protein expression in KG-1a cells

Although bisdemethoxycurcumin demonstrated potential cancer-prevention properties, it was excluded from subsequent experiments due to low SI, suggesting it had limited specificity for KG-1a cells and diminished therapeutic potential. After treatments with IC_{20} concentrations of the crude ethanolic extract of *Z. officinale*, shogaol, gingerol, the crude ethanolic extract of *C. longa*,



Fig. 3 Inhibitory effects of crude ethanolic extracts and their active compound on IL-2 production. PHA-activated PBMCs were incubated with crude ethanolic extracts and their active compounds for 24 h. The supernatants were collected and measured the levels of IL-2 by ELISA kit. All compounds and extracts inhibited IL-2 release by more than 20% compared to dexamethasone (positive control). Data are the mean \pm SD (n=3); *p < 0.05 vs. positive control

curcumin, the crude ethanolic extract of *C. zedoaria*, and demethoxycurcumin, (Table S9) the result showed that WT1 protein levels were significantly decreased by 53.40 ± 6.21 , 30.71 ± 3.90 , 43.98 ± 8.93 , 61.50 ± 3.11 , 44.05 ± 9.81 , 48.33 ± 4.62 , and $33.37\pm3.87\%$, respectively when compared to VC (Figs. 4A and B).

Additionally, the CD34 protein levels after treatment with the crude ethanolic extract of *Z. officinale*, shogaol, gingerol, the crude ethanolic extract of *C. longa*, curcumin, the crude ethanolic extract of *C. zedoaria*, and demethoxycurcumin at IC_{20} concentrations (Table S9) were significantly decreased by 33.70 ± 4.21 , 35.93 ± 3.70 , 31.95 ± 5.70 , 37.99 ± 1.80 , 29.29 ± 2.43 , 28.24 ± 8.31 , and $32.79 \pm 1.56\%$, respectively, compared to VC (Figs. 4A and B).

Interestingly, the results also revealed that all active compounds exhibited a more potent inhibitory effect on WT1 protein expression than their respective crude ethanolic extracts. However, there was no significant difference in the inhibitory effect between the crude ethanolic extracts and their active compounds on CD34 protein levels.

Effect of crude ethanolic extracts and active compounds on total cell number in KG-1a cells

The result showed that the total cell number after treatments with crude ethanolic extract of *Z. officinale*, shogaol, gingerol, crude ethanolic extract of *C. longa*, curcumin, crude ethanolic extract of *C. zedoaria*, and bisdemethoxycurcumin with IC_{20} concentrations (Table S9) significantly decreased by 35.95 ± 12.61 , 35.32 ± 6.10 , 31.39 ± 8.69 , 36.32 ± 6.88 , 40.75 ± 12.78 , 35.97 ± 12.01 , and $47.21 \pm 7.98\%$, respectively, when compared to VC. There is no difference in the inhibitory effect between crude ethanolic extracts and their active compounds on total cell numbers (Fig. 5).

To compare the viable and dead cell proportions across treatment groups, normalized cell counts ($\times 10^3$ cells per 1 million total cells) were calculated. After treatment with the crude ethanolic extracts and their active compounds, the normalized viable cell counts in VC, crude ethanolic extract of Z. officinale, shogaol, gingerol, crude ethanolic extract of C. longa, curcumin, crude ethanolic extract of C. zedoaria, and bisdemethoxycurcumin were $992.15 \pm 7.12, 960.41 \pm 11.10, 976.13 \pm 13.69, 979.50 \pm 8.90,$ 972.17 ± 6.19 , 954.40 ± 3.19 , 961.98 ± 23.65, and $962.81 \pm 19.30 \times 10^3$ cells per 1 million total cells, respectively. The normalized dead cell counts in each treatment group were less than 65×10^3 cells per 1 million total cells, with no significant differences in the proportions of viable and dead cells among treatments (Fig. S9).

Effect of curcumin and shogaol on cell cycle in KG-1a cells

Based on their higher efficacy and SI observed in previous experiments, curcumin and shogaol were prioritized for further evaluation. These compounds significantly reduced WT1 protein expression, supporting their potential as effective anti-leukemic agents. After treatments with various concentrations of active compounds and nocodazole (positive control) in KG-1a cells for 48 h, distinct effects of compounds compared to VC were observed.

Apoptosis was evaluated by measuring the sub- G_1 population using the Nicoletti assay. The results showed that,



Fig. 4 Effect of crude ethanolic extracts and their active compounds treatments on CD34 and WT1 protein expressions in KG-1a cells. KG-1a cells were treated with crude ethanolic extracts and their active compounds with IC_{20} concentrations for 48 h. CD34 and WT1 protein levels were determined by Western blotting. (**A**) Representative images and (**B**) representative bar graph of inhibitory effects of crude ethanolic extracts and their active compounds on WT1 and CD34 protein expressions in KG-1a cells for 48 h. The levels of protein were normalized using GAPDH protein levels. Full-length blots are presented in Supplementary file (Figs. S14–S16). All compounds and extracts inhibited WT1 and CD34 expressions significantly by more than 20% compared to VC. Data are the mean \pm SD (n=3); *p < 0.05 vs. VC



Fig. 5 Antiproliferative effects of crude ethanolic extracts and their active compound in KG-1a cells. KG-1a cells were incubated with various concentrations of crude ethanolic extracts and their active compounds for 48 h. Cells were collected and analyzed by trypan blue exclusion method. All treatments significantly decreased the total cell number compared to VC, with no significant difference between crude ethanolic extracts and active compounds. Data are the mean \pm SD (n=3); *p < 0.05 vs. VC

at its IC₂₀ concentration, 6 ng/mL of nocodazole significantly increased the sub-G₁ population, indicating pronounced apoptosis. In contrast, at their respective IC₂₀ concentrations, 3 μ g/mL of curcumin and 0.12 μ g/mL of shogaol did not significantly increase the sub-G₁ population compared to VC. However, at higher concentrations, both curcumin and shogaol noticeably induced cell death, with significant increases in the sub-G₁ population observed (Figs. 6A–B).

To evaluate the effects of curcumin and shogaol on cell cycle arrest, the sub- G_1 population was excluded, and the distribution of cells across cell cycle phases was analyzed. The results revealed that at the IC₂₀ concentration, nocodazole significantly increased the proportion of cells in the G_2/M phase by $23.48 \pm 1.35\%$, resulting in a significant decrease in the S phase population. In contrast, at their respective IC₂₀ concentrations, neither curcumin nor shogaol induced significant changes in the cell cycle phase population compared to VC. However, 1.5 µg/mL of shogaol significantly decreased the S phase population by $40.56 \pm 3.82\%$, although shogaol did not induce cell cycle arrest. At IC₅₀ concentrations, both curcumin

and shogaol increased the G_2/M phase population to $43.66 \pm 6.75\%$ and $31.38 \pm 1.60\%$, respectively, compared to VC, resulting in a significant decrease in the S phase population (Figs. 6C–D).

Effect of curcumin and shogaol on cell death in KG-1a cells

After KG-1a cells were treated with compounds at their IC_{50} concentrations, the results indicated that both compounds could induce cell death in a dose- and time-dependent manner. Nocodazole (positive control) at concentrations of 5–120 ng/mL showed a significant induction of cell death after 24 h of incubation when compared to VC, and this effect dramatically increased at 48 h of incubation (Fig. 7A). Shogaol at 6–9 µg/mL notably induced cell death at 6 h, and its effect increased continuously across all concentrations with longer incubation times. In contrast, curcumin gradually induced cell death after 12 h of incubation (Figs. 7B and C).



Fig. 6 Effect of curcumin, shogaol, and nocodazole on cell cycle distribution in KG-1a cells. Cells were treated with various concentrations of compounds and nocodazole (positive control) for 48 h, and cell cycle distribution was analyzed by flow cytometry. Cell cycle was analyzed by flow cytometry (**A**) Histogram and (**B**) bar graph of the sub-G₁ population, showing that nocodazole at IC₂₀ significantly increased apoptosis. While curcumin and shogaol at IC₂₀ did not significantly affect the sub-G₁ population, both compounds increased apoptosis at higher concentrations. (**C**) Dot plot and (**D**) bar graph of cell cycle distribution at IC₂₀ but increased the G₂/M phase population and decreased the S phase population at their highest concentrations (IC₅₀). Data are the mean ± SD (n=3); *p < 0.05 vs. VC



Fig. 7 Effect of curcumin, shogaol, and nocodazole on cell death in KG-1a cells. KG-1a cells were incubated with various concentrations of curcumin, shogaol, and nocodazole (positive control) for different incubation times. After collecting the cells, cells were stained with PI solution and analyzed by flow cytometry. Representative bar graph depicting percentage dead cells after treatments with various concentrations and times compared with VC in KG-1a cells. **(A)** Nocodazole induced significant cell death at 24 h, with a dramatic increase at 48 h. **(B)** Curcumin gradually induced cell death, with notable increases after 12 h. **(C)** Shogaol induced cell death, with significant effects at 6 h and further increases over time. Data are the mean \pm SD (n=3); *p < 0.05 vs. VC

Effects of curcumin and shogaol on apoptotic relatedprotein and mechanism of apoptosis in KG-1a cells

To investigate the effects of curcumin and shogaol on apoptotic-related proteins in KG-1a cells, the cells were treated with IC_{50} concentrations of curcumin, shogaol, and nocodazole (positive control) for 0–12 h at 37 °C in a humidified incubator with 5% CO₂.

The results showed that shogaol initiated the activation of cleaved caspase-3 and PARP after 3 h of treatment, whereas curcumin activated both proteins after 6 h. However, nocodazole failed to induce the activation of cleaved caspase-3 and PARP even after 12 h of incubation, compared to VC (Figs. 8A–C).

Furthermore, the levels of WT1 and TAZ protein expressions were investigated using Western blotting after treatments with curcumin and shogaol. The results showed that WT1 protein levels gradually decreased after treatments with curcumin and shogaol compared to non-treatment (0 h incubation time). Notably, the WT1 protein level was significantly decreased by $77.01\pm8.80\%$ after treatments with curcumin for 6 h, and by $22.58\pm3.99\%$ after treatment with shogaol for 3 h when compared to VC. Regarding the TAZ protein, no significant change in its abundance was observed. However, nocodazole failed to reduce WT1 expression even after 12 h of incubation (Figs. 8A, D, and E).

After confirming the presence of apoptosis-related proteins, the mechanism of apoptosis induced by these compounds in KG-1a cells was investigated using Western blotting. To explore this mechanism, the expression levels of the phosphorylated forms of c-Jun (p-c-Jun), SAPK/JNK (p-SAPK/JNK), and Akt (p-Akt (S473)) were examined after treating KG-1a cells with curcumin or shogaol for 0–6 h. The results showed that p-c-Jun and



Fig. 8 Western blot analysis following curcumin, shogaol, and nocodazole treatments in KG-1a cells. KG-1a cells were incubated with IC₅₀ concentration of curcumin, shogaol, and nocodazole (positive control) for 0–12 h. The expression of target proteins was analyzed by Western blotting. (A) Protein levels after treatment with the IC₅₀ concentrations of curcumin (10 µg/mL), shogaol (3 µg/mL), and nocodazole (10 ng/mL) for 0–12 h. (**B**) Protein expressions for cleaved caspase-3, (C) cleaved PARP, (D) WT1, and (E) TAZ in KG-1a cells after 0–12 h of treatment. The levels of protein were normalized using GAPDH protein levels. Full-length blots are presented in Supplementary file (uncropped gels and blot images, Figs. S17–S21). Shogaol induced the activation of cleaved caspase-3 and PARP at 3 h, while curcumin triggered these proteins at 6 h. WT1 expression significantly decreased with both curcumin and shogaol treatments, but nocodazole had no effect. TAZ levels remained unchanged. Data are the mean \pm SD (n = 3); *p < 0.05 vs. VC

p-SAPK/JNK increased after incubation with shogaol or curcumin at 6 h, while p-Akt (S473) decreased at 6 h compared to VC (Fig. S12).

Discussion

Inhibition of LSCs has been a key focus in leukemia treatment so far [37]. Therefore, LSC targeting therapy is a novel strategy aimed at improving patient's quality of life after treatments [38, 39]. In this study, KG-1a cell line was used as an LSC model because it exhibits characteristics similar to those of LSCs, including differentiation, proliferation, and self-renewal [40-42]. Plants from the Zingiberaceae family are widely used in traditional medicine across Southeast Asia due to their accessibility and safety. The exploration of the bioactivities of plants in this family holds promise for drug discovery, particularly as potential anticancer agents. Among the 10 crude ethanolic extracts obtained from the rhizomes of Zingiberaceae plants, each extract exhibited varying cytotoxicity across different cancers and leukemic cell lines. Notably, extracts from C. longa, C. zedoaria, and Z. officinale demonstrated significant cytotoxicity against KG-1a leukemic stem cells. Moreover, these extracts exhibited strong cytotoxic effects on K562, MCF-7, and HeLa cells, with less impact on PBMCs. Therefore, C. longa, C. zedoaria, and Z. officinale were selected as the candidate plants for this study. The active compounds of C. longa and C. zedoaria are curcuminoids. Curcumin is the main compound found in C. longa, whereas demethoxycurcumin is the most abundant in C. zedoaria [43-45]. Z. officinale contains active compounds, namely shogaol and gingerol [46]. The anticancer activities of these active compounds and their crude ethanolic extracts were compared in this study. In the cytotoxicity studies, SI provides insight into the relative safety and efficacy of a compound, assessing its ability to selectively target and impact cancer cells without causing significant harm to normal cells. Values higher than 1 indicate favorable selectivity against cancer cells [35, 36]. In this study, all active compounds exhibited SI values greater than 1, except bisdemethoxycurcumin. Notably, shogaol showed an SI value of 3, indicating its specific targeting ability against KG-1a cells.

Suppressing unregulated inflammatory cytokines (IL-2 and TNF- α) and NO productions from macrophages is one method that can reduce the risk of cancer initiation. In this study, the cancer prevention activities of the active compounds were compared with those of their crude ethanolic extracts. Regarding *C. longa* and *Z. zedoaria* plants, curcuminoids and their crude ethanolic extracts showed no difference in activity. Both compounds decreased the production of IL-2 and TNF- α , which are important inflammatory cytokines as well as NO. Regarding *Z. officinale*, shogaol demonstrated better inhibitory activity against the inflammatory cytokines and NO than either the crude ethanolic extract and gingerol. Therefore, curcuminoids and shogaol emerge as promising compounds for cancer prevention studies.

Previous studies have reported that shogaol, gingerol, and curcuminoids exhibited notable anticancer activity and cancer prevention property. The content of shogaol in the crude ethanolic extract of *Z. officinale* was reported to be very low, constituting less than 2% in fresh rhizome [46]. In the current study, shogaol exhibited significant anticancer activity when compared to gingerol and its crude ethanolic extract. Meanwhile the total curcuminoids content in the crude ethanolic extracts of *C. longa* and *C. zedoaria* is approximately 6% [43, 44]. However, the results indicated no significant difference in cancer prevention and anticancer activities between curcuminoids and their crude ethanolic extracts. After confirming the anticancer activity of the extracts, we further examined the active compounds to investigate their mechanisms of action and clarify their contributions to the overall activity. However, bisdemethoxycurcumin was excluded from further assays due to its low SI, indicating limited specificity toward KG-1a cells and reduced therapeutic potential.

The WT1 protein is a hallmark in leukemia and plays an integral role in leukemogenesis. Increased WT1 protein expression associated with high rates of cell proliferation [47]. In this study, the effect of active compounds compared to their crude ethanolic extracts on WT1 and CD34 expressions in KG-1a cells were investigated. The result showed that at non-cytotoxic doses, all active compounds and their crude ethanolic extracts significantly decreased WT1 protein expression in KG-1a cells, thereby reducing leukemic cell stem cell proliferation. The normalized viable cell counts revealed a consistent decrease in leukemic cell viability across treatments, aligning with the observed WT1 downregulation. This supports the role of WT1 in maintaining leukemic cell proliferation and suggests that the treatments effectively targeted this pathway. Interestingly, each crude ethanolic extract showed a stronger inhibitory effect on WT1 expression than its active compounds. Notably, the crude ethanolic extract of Z. officinale demonstrated greater WT1 downregulation than shogaol, even though shogaol exhibited better cancer prevention properties. This suggests that shogaol may possibly target other leukemiarelated proteins apart from WT1. In this study, CD34, a hallmark of the leukemic stem cells that can be phosphorylated by protein kinase C (PKC), was investigated. The results showed that, although all extracts and active compounds could significantly decreased CD34 expression, the inhibitory effects of each treatment were not significantly different.

Despite extensive research on WT1, few studies have explored the Hippo signaling pathway in leukemia. Previous reports have indicated that YAP protein expression in K562 and HL-60 cell lines plays an important role in cell proliferation in leukemic cells [27, 28]. In contrast, our study revealed that KG-1a leukemic stem cells and other leukemic cell lines (K562 and EoL-1) did not express YAP protein, while TAZ was expressed at a low level in KG-1a cells (Figs. S13). However, no significant changes in TAZ expression were observed upon extract treatments. Although YAP/TAZ have been shown to play important roles in chemoresistance in solid tumors, they may not be as relevant in leukemic stem cell chemoresistance.

Since curcumin and shogaol demonstrated higher efficacy and SI in cytotoxicity studies, both compounds were prioritized for further investigation. They significantly inhibited WT1 protein expression, suggesting their strong potential as anti-leukemic agents. Previous reports have indicated that curcuminoids, particularly curcumin, exhibited antiproliferative effects in various leukemic cell lines and patients' leukemic cells [48, 49]. Moreover, previous studies have shown that curcumin treatment in K562 cells leads to the attenuation of WT1 auto-regulatory function. This occurs through the inhibition of PKCa and c-Jun N-terminal kinase (JNK) signaling, which suppresses both WT1 auto-regulation and c-Jun/AP-1 binding to its cognate consensus site at the proximal WT1 gene promoter [50, 51]. Furthermore, curcumin decreased cell proliferation by reducing WT1 protein levels in KG-1a stem cell lines [32]. Meanwhile, 6-shogaol from Z. officinale played a major role in inducing apoptosis in acute lymphoblastic leukemia (ALL) by activating p53 and generating oxidative stress [52]. Additionally, shogaol suppressed proliferation in oral squamous cell carcinoma (OSCC) cells and induced apoptosis by inhibiting the phosphatidylinositol 3-kinase (PI3K)/ Akt signaling pathway, while regulating apoptosis-associated factors such as p53, Bax, Bcl-2, and cleaved caspase-3 [53]. This study is the first to report that shogaol can inhibit WT1 and CD34 expressions.

The PI3K/Akt signaling pathway and JNK pathway regulate numerous cellular processes, including cell growth, migration, invasion, and apoptosis. Phosphorylation of Akt enhances CDK2 activity, which is required for cell progression from the S to G_2/M phase and inhibits apoptosis [54-56]. Therefore, the inhibition of PI3K/ Akt can induce caspase-dependent apoptosis [55, 57]. In this study, curcumin and shogaol were selected for further investigation due to their efficacy as active compounds from their respective candidate plants. While both curcumin and shogaol demonstrated the ability to decrease WT1 protein levels and inhibit cell proliferation at the IC₂₀ concentrations, non-cytotoxic doses were insufficient to arrest the cell cycle in KG-1a cells. IC₅₀ concentrations were required to induce cell cycle arrest at the G_2/M phase and promote cell death. Moreover, time-course experiments indicated that both compounds could induce cell death in a dose- and time-dependent manner. To explore the mechanism of curcumin and shogaol in inducing cell death in KG-1a cells, the cells were treated with the IC₅₀ concentration of each compound. The results indicated that both curcumin and shogaol induced apoptosis by decreasing the phosphorylation of Akt (S473), leading to an increase in the levels of phosphorylated JNK and c-Jun. Subsequently, this process resulted in an upregulation of cleaved caspase-3 and PARP proteins. Additionally, both curcumin and shogaol decreased the expression of WT1 and TAZ proteins, which are associated with leukemic cell proliferation.

Conclusion

Our findings suggest that *C. longa, C. zedoaria*, and *Z. officinale* extracts are potential candidates for cancer treatment, exhibiting significant cytotoxicity across various cancer cell types. Additionally, these extracts and their active compounds demonstrate cancer prevention properties by inhibiting the production of IL-2, TNF- α , and NO. Furthermore, curcumin and shogaol can arrest the cell cycle and induce apoptosis in KG-1a cells through the inhibition of the Akt pathway. Thus, curcumin and shogaol emerge as promising compounds for leukemia treatment, requiring further investigation.

Abbreviations

Abbieviations	
ALL	Acute lymphoblastic leukemia cells
AML	Acute myeloid leukemia
CDK2	Cyclin-dependent kinase 2
CML	Chronic myeloid leukemia
DMEM	Dulbecco's Modified Eagle Medium
HSCs	Hematopoietic stem cells
IC ₂₀	Non-cytotoxic dose (Inhibitory concentration values at 20%
	growth)
IC ₅₀	Inhibitory concentration values at 50% growth
IL-2	Interleukin-2
IMDM	Iscove's Modified Dulbecco's Medium
JNK	c-Jun N-terminal kinase
LPS	Lipopolysaccharide
LSCs	Leukemic stem cells
MRD	Minimal residual disease
NCI	National Cancer Institute
NO	Nitric oxide
OD	Optical density
OSCC	Oral squamous cell carcinoma
p-c-Jun	Phosphorylated forms of c-Jun
p-SAPK/JNK	Phosphorylated forms of SAPK/JNK
p-Akt	Phosphorylated form of Akt
PBMCs	Peripheral blood mononuclear cells
PBS	Phosphate buffer saline
PBS-T	Phosphate buffer saline with Tween-20
PHA	Phytohemagglutinin
PI	Propidium iodide
PI3K	Phosphatidylinositol 3-kinase
PKC	Protein kinase C
RPMI-1640	Roswell Park Memorial Institute-1640
RT	Room temperature
SD	Standard derivation
SI	Selectivity index
TNF-a	Tumor necrosis factor-alpha
VC	Vehicle control
WT1	Wilms' tumor 1

Supplementary Information

The online version contains supplementary material available at https://doi.or g/10.1186/s12906-025-04829-7.

Supplementary Material 1

Supplementary Material 2

Acknowledgements

The authors would like to thank the Department of Medical Technology, Faculty of Associated Medical Sciences and the Center of Excellence in Pharmaceutical Nanotechnology, Chiang Mai University for their equipment and facility support. Additionally, thanks are extended to the Department of Molecular and Medical Pharmacology, Faculty of Life Sciences, Kumamoto University, for their valuable contributions.

Author contributions

Conceptualization: SO, SA, MT. Methodology: SA, PP, SC, ST, YS, MT. Data curation and formal analysis: SA, PP, SC, ST, NV. Resources: SO, SA, MT. Writing-original draft: PP, SO, SA. Writing- review & editing: SO, SA, YS, MT.

Funding

This work was supported by a grant from the Fundamental Fund (180072_FF66/046), Chiang Mai University, Chiang Mai, Thailand and the Agricultural Research Development Agency, Thailand (Grant No. GSCMU(HRD6405069)/08/2564).

Data availability

The datasets used and/or analyzed during the current study are available from the corresponding author by reasonable request.

Declarations

Ethics approval and consent to participate

The use of human PBMCs in this study was approved by the Human Research Ethics Unit of the Faculty of Associated Medical Sciences, Chiang Mai University. The approval numbers were AMSEC-64EX-011 and AMSEC-66EX-117. Written informed consent was obtained from all healthy subjects included in the study.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Medical Technology, Faculty of Associated Medical Sciences, Chiang Mai University, Chiang Mai 50200, Thailand

²Cancer Research Unit of Associated Medical Sciences (AMS CRU), Faculty of Associated Medical Sciences, Chiang Mai University, Chiang Mai 50200, Thailand

³Center of Excellence in Pharmaceutical Nanotechnology, Chiang Mai University, Chiang Mai 50200, Thailand

⁴Center for Metabolic Regulation of Healthy Aging, Faculty of Life Sciences, Kumamoto University, Kumamoto 860-8556, Japan ⁵Department of Pharmaceutical Sciences, Faculty of Pharmacy, Chiang

Mai University, Chiang Mai 50200, Thailand

⁶Department of Molecular and Medical Pharmacology, Faculty of Life Sciences, Kumamoto University, Kumamoto 860-8556, Japan

Received: 29 June 2024 / Accepted: 13 February 2025 Published online: 28 February 2025

References

- Danciu C, Vlaia L, Fetea F, Hancianu M, Coricovac DE, Ciurlea SA, Şoica CM, Marincu I, Vlaia V, Dehelean CA, et al. Evaluation of phenolic profile, antioxidant and anticancer potential of two main representants of Zingiberaceae family against B164A5 murine melanoma cells. Biol Res. 2015;48(1):1. https:// doi.org/10.1186/0717-6287-48-1.
- Manna I, Das D, Mondal S, Bandyopadhyay M. Potential pharmacotherapeutic phytochemicals from Zingiberaceae for cancer prevention. In: Pharmacotherapeutic Botanicals for Cancer Chemoprevention. edn. Edited by Kumar M, Sharma A, Kumar P. Singapore: Springer; 2020; 221–81. https://doi.org/10. 1007/978-981-15-5999-0_10
- Mahomoodally MF, Aumeeruddy MZ, Rengasamy KRR, Roshan S, Hammad S, Pandohee J, Hu X, Zengin G. Ginger and its active compounds in cancer therapy: from folk uses to nano-therapeutic applications. Semin Cancer Biol. 2021;69:140–9. https://doi.org/10.1016/j.semcancer.2019.08.009.
- Ballester P, Cerdá B, Arcusa R, García-Muñoz AM, Marhuenda J, Zafrilla P. Antioxidant activity in extracts from Zingiberaceae family: cardamom, turmeric, and ginger. Molecules. 2023;28(10):4024. https://doi.org/10.3390/molecules2 8104024.
- Coussens LM, Werb Z. Inflammation and cancer. Nature. 2002;420(6917):860– 7. https://doi.org/10.1038/nature01322.

- Greten FR, Grivennikov SI. Inflammation and cancer: triggers, mechanisms, and consequences. Immunity. 2019;51(1):27–41. https://doi.org/10.1016/j.im muni.2019.06.025.
- Zhao H, Wu L, Yan G, Chen Y, Zhou M, Wu Y, Li Y. Inflammation and tumor progression: signaling pathways and targeted intervention. Signal Transduct Target Therapy. 2021;6(1):263.
- Ko J-H, Sethi G, Um J-Y, Shanmugam MK, Arfuso F, Kumar AP, Bishayee A, Ahn KS. The role of resveratrol in cancer therapy. Int J Mol Sci. 2017;18(12):2589. ht tps://doi.org/10.3390/ijms18122589.
- Ruhee RT, Ma S, Suzuki K. Sulforaphane protects cells against lipopolysaccharide-stimulated inflammation in murine macrophages. Antioxid (Basel). 2019;8(12):577. https://doi.org/10.3390/antiox8120577.
- Makon-Sébastien N, Francis F, Eric S, Henri VP, François LJ, Laurent P, Yves B, Serge C. Lycopene modulates THP1 and Caco2 cells inflammatory state through transcriptional and nontranscriptional processes. Mediators Inflamm. 2014;2014:507272. https://doi.org/10.1155/2014/507272.
- Mansouri K, Rasoulpoor S, Daneshkhah A, Abolfathi S, Salari N, Mohammadi M, Rasoulpoor S, Shabani S. Clinical effects of curcumin in enhancing cancer therapy: a systematic review. BMC Cancer. 2020;20(1):791. https://doi.org/10.1 186/s12885-020-07256-8.
- Jia Y, Li X, Meng X, Lei J, Xia Y, Yu L. Anticancer perspective of 6-shogaol: anticancer properties, mechanism of action, synergism and delivery system. Chin Med. 2023;18(1):138. https://doi.org/10.1186/s13020-023-00839-0.
- Todoric J, Antonucci L, Karin M. Targeting inflammation in cancer prevention and therapy. Cancer Prev Res (Phila). 2016;9(12):895–905. https://doi.org/10.1 158/1940-6207.Capr-16-0209.
- Dick JE. Acute myeloid leukemia stem cells. Ann N Y Acad Sci. 2005;1044(1):1– 5. https://doi.org/10.1196/annals.1349.001.
- Hanekamp D, Cloos J, Schuurhuis GJ. Leukemic stem cells: identification and clinical application. Int J Hematol. 2017;105(5):549–57. https://doi.org/10.100 7/s12185-017-2221-5.
- Fraizer G, Patmasiriwat P, Zhang X, Saunders G. Expression of the tumor suppressor gene WT1 in both human and mouse bone marrow [letter]. Blood. 1995;86(12):4704–6. http://www.bloodjournal.org/content/bloodjournal/86/ 12/4704.full.pdf.
- 17. Baird PN, Simmons PJ. Expression of the *Wilms' tumor* gene (*WT1*) in normal hemopoiesis. Exp Hematol. 1997;25(4):312–20.
- Maurer U, Weidmann E, Karakas T, Hoelzer D, Bergmann L. Wilms tumor gene (WT1) mRNA is equally expressed in blast cells from acute myeloid leukemia and normal CD34⁺ progenitors. Blood. 1997;90(10):4230–2. https://doi.org/10 .1182/blood.V90.10.4230.
- Maurer U, Brieger J, Weidmann E, Mitrou P, Hoelzer D, Bergmann L. The Wilms' tumor gene is expressed in a subset of CD34⁺ progenitors and downregulated early in the course of differentiation *in vitro*. Exp Hematol. 1997;25(9):945–50.
- Hosen N, Sonoda Y, Oji Y, Kimura T, Minamiguchi H, Tamaki H, Kawakami M, Asada M, Kanato K, Motomura M. Very low frequencies of human normal CD34⁺ haematopoietic progenitor cells express the Wilms' tumour gene WT1 at levels similar to those in leukaemia cells. Br J Haematol. 2002;116(2):409–20.
- 21. Lazzarotto D, Candoni A. The role of *Wilms' tumor* gene (*WT1*) expression as a marker of minimal residual disease in acute myeloid leukemia. J Clin Med. 2022;11(12):3306. https://doi.org/10.3390/jcm11123306.
- Bergmann L, Miething C, Maurer U, Brieger J, Karakas T, Weidmann E, Hoelzer D. High levels of *Wilms' tumor* gene (*WT1*) mRNA in acute myeloid leukemias are associated with a worse long-term outcome. Blood. 1997;90(3):1217–25.
- Boublikova L, Kalinova M, Ryan J, Quinn F, O'Marcaigh A, Smith O, Browne P, Stary J, McCann SR, Trka J et al. *Wilms' tumor* gene 1 (*WT1*) expression in childhood acute lymphoblastic leukemia: a wide range of WT1 expression levels, its impact on prognosis and minimal residual disease monitoring. Leukemia. 2005;20(2):254–63. https://doi.org/10.1038/sj.leu.2404047
- 24. Tamaki H, Ogawa H, Inoue K, Soma T, Yamagami T, Miyake S, Oka Y, Oji Y, Tatekawa T, Tsuboi A, et al. Increased expression of the Wilms tumor gene (WT1) at relapse in acute leukemia [letter]. Blood. 1996;88(11):4396–8. http:// www.bloodjournal.org/content/bloodjournal/88/11/4396.2.full.pdf.
- Harvey KF, Zhang X, Thomas DM. The Hippo pathway and human cancer. Nat Rev Cancer. 2013;13(4):246–57. https://doi.org/10.1038/nrc3458.
- Calses PC, Crawford JJ, Lill JR, Dey A. Hippo pathway in cancer: aberrant regulation and therapeutic opportunities. Trends Cancer. 2019;5(5):297–307. https://doi.org/10.1016/j.trecan.2019.04.001.
- 27. Li H, Huang Z, Gao M, Huang N, Luo Z, Shen H, Wang X, Wang T, Hu J, Feng W. Inhibition of YAP suppresses CML cell proliferation and enhances efficacy of

imatinib *in vitro* and *in vivo*. J Exp Clin Cancer Res. 2016;35(1):134. https://doi. org/10.1186/s13046-016-0414-z.

- Chen M, Wang J, Yao S-F, Zhao Y, Liu L, Li L-W, Xu T, Gan L-G, Xiao C-L, Shan Z-L, et al. Effect of YAP inhibition on human leukemia HL-60 cells. Int J Med Sci. 2017;14(9):902–10. https://doi.org/10.7150/ijms.19965.
- 29. Tachakittirungrod S, Okonogi S, Chowwanapoonpohn S. Study on antioxidant activity of certain plants in Thailand: mechanism of antioxidant action of guava leaf extract. Food Chem. 2007;103(2):381–8.
- Limtrakul P, Anuchapreeda S, Buddhasukh D. Modulation of human multidrug-resistance MDR-1 gene by natural curcuminoids. BMC Cancer. 2004;4(1):13. https://doi.org/10.1186/1471-2407-4-13.
- Chearwae W, Anuchapreeda S, Nandigama K, Ambudkar SV, Limtrakul P. Biochemical mechanism of modulation of human P-glycoprotein (ABCB1) by curcumin I, II, and III purified from turmeric powder. Biochem Pharmacol. 2004;68(10):2043–52://WOS:000224714300015.
- Panyajai P, Tima S, Chiampanichayakul S, Anuchapreeda S. Dietary turmeric bisdemethoxycurcumin suppresses Wilms' tumor 1 and CD34 protein expressions in KG-1a leukemic stem cells. Nutr Cancer. 2019;71(7):1189–200. https:// doi.org/10.1080/01635581.2019.1598565.
- Omoruyi SI, Kangwa TS, Ibrakaw AS, Cupido CN, Marnewick JL, Ekpo OE, Hussein AA. Cytotoxic activities of selected plants of the family Amaryllidaceae on brain tumour cell lines. S Afr J Bot. 2021;136:118–25. https://doi.org/10.10 16/j.sajb.2020.09.015.
- Suffness M, John M. Pezzuto; Assays related to cancer drug discovery. In: Methods in Plant Biochemistry: Assays for Bioactivity. 6th edn. Edited by Hostettmann K. London, UK: Academic Press; 1990: 33–71.
- Indrayanto G, Putra GS, Suhud F. Chapter six Validation of *in-vitro* bioassay methods: application in herbal drug research. In: Profiles of Drug Substances, Excipients and Related Methodology. Edited by Al-Majed AA, vol. 46. USA: Academic Press; 2021: 273–307. https://doi.org/10.1016/bs.podrm.2020.07.00
- Lica JJ, Wieczór M, Grabe GJ, Heldt M, Jancz M, Misiak M, Gucwa K, Brankiewicz W, Maciejewska N, Stupak A, et al. Effective drug concentration and selectivity depends on fraction of primitive cells. Int J Mol Sci. 2021;22(9):4931. https://doi.org/10.3390/ijms22094931.
- Rodrigues ACBC, Costa RGA, Silva SLR, Dias IRSB, Dias RB, Bezerra DP. Cell signaling pathways as molecular targets to eliminate AML stem cells. Crit Rev Oncol Hematol. 2021;160:103277. https://doi.org/10.1016/j.critrevonc.2021.1 03277.
- Valent P, Sadovnik I, Eisenwort G, Bauer K, Herrmann H, Gleixner KV, Schulenburg A, Rabitsch W, Sperr WR, Wolf D. Immunotherapy-based targeting and elimination of leukemic stem cells in AML and CML. Int J Mol Sci. 2019;20(17):4233. https://doi.org/10.3390/ijms20174233.
- Khaldoyanidi SK, Hindoyan A, Stein A, Subklewe M. Leukemic stem cells as a target for eliminating acute myeloid leukemia: gaps in translational research. Crit Rev Oncol Hematol. 2022;175:103710. https://doi.org/10.1016/j.critrevon c.2022.103710.
- Fuchs D, Daniel V, Sadeghi M, Opelz G, Naujokat C. Salinomycin overcomes ABC transporter-mediated multidrug and apoptosis resistance in human leukemia stem cell-like KG-1a cells. Biochem Biophys Res Commun. 2010;394(4):1098–104. https://doi.org/10.1016/j.bbrc.2010.03.138.
- She M, Niu X, Chen X, Li J, Zhou M, He Y, Le Y, Guo K. Resistance of leukemic stem-like cells in AML cell line KG1a to natural killer cell-mediated cytotoxicity. Cancer Lett. 2012;318(2):173–9.
- 42. O'Reilly E, Zeinabad HA, Nolan C, Sefy J, Williams T, Tarunina M, Hernandez D, Choo Y, Szegezdi E. Recreating the bone marrow microenvironment to model leukemic stem cell quiescence. Front Cell Dev Biol. 2021;9:662868. htt ps://doi.org/10.3389/fcell.2021.662868.
- Paramapojn S, Gritsanapan W. Variation of curcuminoids in ethanolic extract of *Curcuma zedoaria* rhizomes in Thailand by HPLC. Planta Med. 2007;73(09):227. https://doi.org/10.1055/s-2007-987008.

- Lobo R, Prabhu KS, Shirwaikar A, Shirwaikar A. *Curcuma zedoaria* Rosc. (white turmeric): a review of its chemical, pharmacological and ethnomedicinal properties. J Pharm Pharmacol. 2009;61(1):13–21. https://doi.org/10.1211/jpp. 61.01.0003.
- Kocaadam B, Şanlier N. Curcumin, an active component of turmeric (*Curcuma longa*), and its effects on health. Crit Rev Food Sci Nutr. 2017;57(13):2889–95. https://doi.org/10.1080/10408398.2015.1077195.
- Seon O, Woo-Sik J. Optimization of extraction conditions for the 6-shogaolrich extract from ginger (*Zingiber officinale* Roscoe). Prev Nutr Food Sci. 2012;17(2):166–71. https://doi.org/10.3746/pnf.2012.17.2.166.
- Zhou B, Jin X, Jin W, Huang X, Wu Y, Li H, Zhu W, Qin X, Ye H, Gao S. WT1 facilitates the self-renewal of leukemia-initiating cells through the upregulation of BCL2L2: WT1-BCL2L2 axis as a new acute myeloid leukemia therapy target. J Transl Med. 2020;18(1):254. https://doi.org/10.1186/s12967-020-02384-y.
- Anuchapreeda S, Limtrakul P, Thanarattanakorn P, Sittipreechacharn S, Chanarat P. Inhibitory effect of curcumin on WT1 gene expression in patient leukemic cells. Arch Pharm Res. 2006;29(1):80–7. https://doi.org/10.1007/BF0 2977473.
- Anuchapreeda S, Tima S, Duangrat C, Limtrakul P. Effect of pure curcumin, demethoxycurcumin, and bisdemethoxycurcumin on WT1 gene expression in leukemic cell lines. Cancer Chemother Pharmacol. 2008;62(4):585–94. https ://doi.org/10.1007/s00280-007-0642-1.
- Semsri S, Krig SR, Kotelawala L, Sweeney CA, Anuchapreeda S. Inhibitory mechanism of pure curcumin on Wilms' tumor 1 (WT1) gene expression through the PKCa signaling pathway in leukemic K562 cells. FEBS Lett. 2011;585(14):2235–42. http://www.sciencedirect.com/science/article/pii/S00 14579311004091.
- Anuchapreeda S, Rungrojsakul M, Tima S, Chiampanichayakul S, Krig SR. Coactivation of WT1 and AP-1 proteins on WT1 gene promoter to induce WT1 gene expression in K562 cells. Cell Signal. 2019;53:339–47. https://doi.org/10. 1016/j.cellsig.2018.11.001.
- 52. Najafi Dorcheh S, Rahgozar S, Talei D. 6-Shogaol induces apoptosis in acute lymphoblastic leukaemia cells by targeting p53 signalling pathway and generation of reactive oxygen species. J Cell Mol Med. 2021;25(13):6148–60. https://doi.org/10.1111/jcmm.16528.
- Huang H, Kim M-O, Kim K-R. Anticancer effects of 6-shogaol via the AKT signaling pathway in oral squamous cell carcinoma. J Appl Oral Sci. 2021;29. https://doi.org/10.1590/1678-7757-2021-0209.
- Maddika S, Ande SR, Wiechec E, Hansen LL, Wesselborg S, Los M. Akt-mediated phosphorylation of CDK2 regulates its dual role in cell cycle progression and apoptosis. J Cell Sci. 2008;121(7):979–88. https://doi.org/10.1242/jcs.0095 30.
- Zhao H-F, Wang J, Tony To S-S. The phosphatidylinositol 3-kinase/Akt and c-Jun N-terminal kinase signaling in cancer: alliance or contradiction? (review). Int J Oncol. 2015;47(2):429–36. https://doi.org/10.3892/ijo.2015.305
- Tsai P-J, Lai Y-H, Manne RK, Tsai Y-S, Sarbassov D, Lin H-K. Akt: a key transducer in cancer. J Biomed Sci. 2022;29(1):76. https://doi.org/10.1186/s12929-022-00 860-9.
- Jeong S-J, Dasgupta A, Jung K-J, Um J-H, Burke A, Park HU, Brady JN. PI3K/AKT inhibition induces caspase-dependent apoptosis in HTLV-1-transformed cells. Virology. 2008;370(2):264–72. https://doi.org/10.1016/j.virol.2007.09.003.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.